

Research article

Human glutaminyl cyclase and bacterial zinc aminopeptidase share a common fold and active site

Rachell E Booth¹, Simon C Lovell², Stephanie A Misquitta³ and Robert C Bateman Jr^{*3}

Address: ¹Department of Chemistry and Biochemistry, Texas State University-San Marcos, San Marcos, TX 78666, USA, ²School of Biological Sciences, University of Manchester, Manchester, M13 9PT, UK and ³Department of Chemistry and Biochemistry, University of Southern Mississippi, Hattiesburg, MS 39406-5043, USA

Email: Rachell E Booth - rbooth@swt.edu; Simon C Lovell - simon.lovell@man.ac.uk; Stephanie A Misquitta - stephaniemisquitta@yahoo.com; Robert C Bateman* - robert.bateman@usm.edu

* Corresponding author

Published: 10 February 2004

Received: 04 November 2003

BMC Biology 2004, 2:2

Accepted: 10 February 2004

This article is available from: <http://www.biomedcentral.com/1741-7007/2/2>

© 2004 Booth et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Glutaminyl cyclase (QC) forms the pyroglutamyl residue at the amino terminus of numerous secretory peptides and proteins. We previously proposed the mammalian QC has some features in common with zinc aminopeptidases. We now have generated a structural model for human QC based on the aminopeptidase fold (pdb code IAMP) and mutated the apparent active site residues to assess their role in QC catalysis.

Results: The structural model proposed here for human QC, deposited in the protein databank as IMOI, is supported by a variety of fold prediction programs, by the circular dichroism spectrum, and by the presence of the disulfide. Mutagenesis of the six active site residues present in both IAMP and QC reveal essential roles for the two histidines (140 and 330, QC numbering) and the two glutamates (201 and 202), while the two aspartates (159 and 248) appear to play no catalytic role. ICP-MS analysis shows less than stoichiometric zinc (0.3:1) in the purified enzyme.

Conclusions: We conclude that human pituitary glutaminyl cyclase and bacterial zinc aminopeptidase share a common fold and active site residues. In contrast to the aminopeptidase, however, QC does not appear to require zinc for enzymatic activity.

Background

The synthesis of biologically active peptide hormones and neurotransmitters begins with precursor proteins undergoing maturation via different processing enzymes. Pro-hormone convertase catalyzes cleavage within precursor proteins at paired basic amino acid residues [1] while carboxypeptidase removes these basic residues [2]. Several peptide hormones and neurotransmitters, such as thyrotropin-releasing hormone, are further modified at their termini. These termini modifications appear to be neces-

sary for recognition of a receptor-binding site or protection of the peptide from exopeptidases. A frequent terminus modification is the oxidative cleavage by peptidyl α -amidating monooxygenase [3] of a carboxyl terminal glycine to form a terminal amide. The amino terminus can be acylated or, in the case of an N-terminal glutaminyl residue, can be cyclized to the pyroglutamyl residue with the accompanying release of the amide nitrogen as ammonia. This last reaction is catalyzed by glutaminyl cyclase (EC 2.3.2.5).

In 1965 Messer and Ottesen identified and purified the first known form of glutaminyl cyclase (QC) from papaya latex [4]. The laboratories of J.S. Kizer and J. Spiess concurrently published the initial identification of QC in porcine and bovine pituitaries, respectively [5,6] and, as expected, QC was found to be localized within secretory vesicles [7]. Subsequent cloning of the bovine and human QC cDNA revealed very similar proteins of 361 amino acids, including a nearly thirty amino acid endoplasmic reticulum signal sequence [8,9]. QC homologs can be discerned in the genomes of a wide variety of organisms including yeast [10], with the human QC gene residing on chromosome 2 (NCBI LocusID: 25797).

Human pituitary QC has been expressed in bacteria, although this has often resulted in protein with little activity, stability, or solubility, or active protein with a non-cleavable GST fusion [9-12]. Thus, functional and structural characterization of QC has been limited. Site directed mutagenesis of the two conserved cysteines revealed neither was essential to enzyme function [12]. Recent mutational analysis of four highly conserved histidine residues indicated that two residues participated in catalysis and two in substrate binding [10]. Taken together with the apparent lack of nucleophilic essential amino acids [5,12], this suggests catalysis by mammalian QC proceeds via general acid/base promotion of the direct cyclization of the terminal glutamine as we have proposed for the papaya enzyme [13,14].

Glutaminyl cyclase has also been purified from papaya latex, but animal and plant forms of QC have no discernable primary sequence identity [15]. We hypothesized previously [10] that, while animal and plant forms of QC may utilize similar catalytic mechanisms, they have different tertiary structures and probably represent two independent families of enzymes. Taken together with previous work on papaya QC, this study demonstrates that the mammalian and plant QC forms are indeed structurally unrelated, while providing evidence of a structural relationship between mammalian QC and the bacterial zinc aminopeptidase family.

Results

Features of the model of human QC


We predict that human QC and the bacterial zinc exopeptidases share a similar fold (Figure 1). This consists of a central 8-stranded β -sheet, which is of a mixed parallel/anti-parallel nature. The sheet is flanked by six α -helices on one side and three on the other. The proposed active site lies at the C-terminal end of the central two β -strands, and is mostly comprised of residues from loop regions.

The sequence identity between human QC and members of the aminopeptidase family is low: between QC and *Vibrio proteolyticus* aminopeptidase (pdb code 1AMP, [16]), for example, it is 16% over the entire length of the alignment. Nevertheless the fold recognition program "Fugue" assigns QC to the "bacterial exopeptidases" family with a Z-score of 26.5. Z-scores with values above 6 correspond to "certain" assignments (>99% probability). Fugue is amongst the most sensitive fold recognition programs currently available [17]. It considers not just the amino acid type, but also its environment, up-weighting those amino-acid residues likely to be conserved through evolution. Conservation of these important features leads Fugue to give the confident assignment in this case.

The sequence alignment (Figure 2) and model have several convincing features. The Pspred prediction of secondary structure corresponds extremely well to that of the aminopeptidase family (Figure 2). This family consists of aminopeptidases from *Vibrio proteolyticus* (pdb code 1AMP) and *Streptomyces griseus* (pdb code 1XJO, [18], and the catalytic domain of carboxypeptidase G2 from *Pseudomonas* (pdb code 1CG2, [19]). Fugue does not use predicted secondary structure to determine either the most likely family assignment for a given sequence nor the alignment, and so the correlation gives increased confidence in the fold assignment and alignment. The N-linked glycosylation sites (Asn 49 and Asn 296) are solvent exposed in the model. The cysteine residues are positioned such that they are able to form a disulfide bond. This bond is in a different position in the QC model as compared to that in template structures 1AMP and 1XJO (the disulfide is absent in 1CG2). In the QC model the disulfide is between α -helix 4 and the third β -strand of the sheet, and as such is completely buried in the core and unlikely to play any direct role in catalysis. The reported effect on activity produced by reducing the disulfide [11] is therefore likely to be structural.

We predict from the model, therefore, that human QC has a mixed α -helix/ β -strand structure that, consistent with the previous prediction, Asn 49 is the likely glycosylation site, and that the two cysteine residues (139 and 164) should be in the disulfide-bonded rather than free thiol form. It should be noted that these features were not used as constraints when aligning the sequences. Their experimental validation can, therefore, serve as a confirmation of the model.

Interestingly, the active site residues (His 140, Asp 159, Glu 201, Glu 202, Asp 248 and His 330, QC numbering) in 1AMP are conserved in the QC model (Figure 3), and probably adopt a similar conformation. Five of these residues form a metal binding site for two zinc ions in 1AMP,

**Figure 1**

Ribbon diagram of hQC theoretical model. Ribbon diagram of secondary and tertiary structures of *Vibrio* aminopeptidase (PDB 1AMP) and human glutaminyl cyclase (PDB 1MOI). Structures are oriented so that active sites are on the right of the molecule. Illustration was generated with DeepView 3.7 using solid open GL rendering and colored by secondary structure succession.

and it would appear that a metal binding site could also be present in human QC. However, there is conflicting evidence for the presence of zinc in human QC (see discussion).

Analysis of papaya QC

In contrast to human QC, the papaya enzyme produces hits with lower confidence levels. It is predicted by Fugue [17] as having similarity to the cytochrome cd1-nitrite reductase family, specifically to the β -sheet domain, which is an eight-bladed β -propeller structure (pdb codes 1AOQ and 1NIR, Z-score = 8.1) and to bacterial amine dehydrogenase (pdb code 1JJU, chain B, Z-score = 7.4), which is a seven-bladed β -propeller. Again in contrast to human QC, papaya QC is predicted by Psipred to have an entirely β -sheet structure, which is consistent with the CD spectrum [20]. Given the relatively low score, the ambiguity regarding the number of blades of the propeller and the difficulty of building accurate models of proteins with repetitive secondary structure [21], we have not built a model of papaya QC. However, it is clear that the papaya QC is structurally distinct from the human QC having an all-beta structure, possibly a β -propeller.


Secondary structure and metal analyses

Secondary structural characteristics of DES hQC and the mutant forms (data not shown) were examined by circular dichroism (Figure 4). Samples were prepared as described in the methods section and then analyzed at Tulane University with an OLIS CD spectrophotometer. Eight scans from 260 nm to 196 nm were obtained on a 6 μ M sample. The resulting spectrum indicated that DES hQC contains 32% α -helix, and 17% β -sheet, a secondary structure composition that correlates well with the *Vibrio proteolyticus* aminopeptidase, which contains 35% α -helix, and 17% β -sheet.

Metal content was determined by ICP-MS as described in Methods. An average of 2.0 nmol/ml Zn were obtained for every 6.2 nmol/ml QC, giving a molar stoichiometry of 0.32:1 Zn:QC.

Active site identification

Four QC acidic amino acid residues were selected for site-directed mutagenesis to alanine based on their conservation between species [10] as well as their presence in the bacterial aminopeptidase active site (see figure 3 and model analysis above). Mutations in either glutamate 201

**Figure 2**

Sequence alignment of hQC and structural templates. Alignment of amino acid sequence of *Vibrio* aminopeptidase (PDB code 1AMP), *Streptomyces* aminopeptidase (PDB code 1XJO) *Pseudomonas* carboxypeptidase G2 (PDB code 1CG2) and human pituitary glutaminyl cyclase (bottom line). Numbers along the top indicate the alignment position. Numbers in parentheses on the left hand side indicate the sequence numbering for the first amino acid in on that line for each sequence. Bars indicate secondary structure (dark bars for helices, light bars for strands) for those sequences with experimentally determined structures. Bars underneath the alignment (labeled "Psipred") indicate predicted secondary structure for the human QC sequence.



**Figure 3****Predicted active site of human glutaminyl cyclase.**

Figure 5 represents the five conserved human QC residues corresponding to the zinc ligands of the bacterial aminopeptidase plus the conserved glutamate corresponding to the general base of the aminopeptidase. QC numbering from the top (clockwise): His 330, Glu 202, Glu 201, His 140, Asp 248, Asp 159. See Table 1 for comparisons with IAMP.

or 202 resulted in completely inactive QC forms. These mutants interact as well as unmutated enzyme with the affinity column, suggesting that the binding site was not disrupted; furthermore, fluorescence scans and CD spectroscopy revealed no apparent changes in structure. Kinetic studies on aspartate 159 and 248 mutants revealed no significant differences in the K_m from unmutated DES hQC (1.4 ± 0.2 (unmutated), 1.5 ± 0.3 (D159), and 1.4 ± 0.4 (D248) mM, respectively) and only a slight decrease in the V_{max} of D159 (3.7 ± 0.4 (unmutated), 1.5 ± 0.2 (D159), and 3.3 ± 0.5 (D248) $\mu\text{mol}/\text{min}/\text{mg}$, respectively).

Discussion

As a peptide-processing enzyme in the mammalian secretory pathway, QC is susceptible to a variety of posttranslational modifications. Recently we showed that at least one of the two predicted N-linked glycosylation sites is indeed glycosylated in the insect-expressed protein and one, if not both, sites in the native tissue QC contain glycans as well [22]. Mutagenesis of both sites showed that the presence or absence of glycans had no apparent effect on QC enzymatic activity. The glycans, then, either have a cellular role such as localization or they are simply fortuitous. Also, consistent with the oxidizing environment of the secretory pathway is the finding that the two cysteine residues in QC are disulfide-bonded [22]. The presence of

**Figure 4**

Circular dichroism spectrum of DES hQC. Circular dichroism of 6 μM DES hQC in 5 mM sodium phosphate buffer, pH 6.0 from 196 to 260 nm.

a disulfide bond was also recently reported by Schilling [11] in human QC expressed in yeast. They postulated a role for this disulfide bond in enzyme activity since addition of high concentrations of reducing agents result in loss of activity. Our previous mutagenesis of these two cysteine residues [12] indicate that they are not directly involved in catalysis, making it likely that they play a role in stabilizing an active form of enzyme tertiary structure. Indeed, the model predicts the disulfide bond is immediately behind the enzyme active site and crosslinks two pieces of secondary structure directly connected to loops containing active site residues.

Tertiary structure examination of QC began with the development of a predicted model (pdb code 1MOI) based on a bacterial zinc aminopeptidase fold as described in the methods section and predicted previously [10]. DES hQC secondary structure composition determined using circular dichroism shows DES hQC to contain 32% α -helix, 17% β -sheet, and 52% random coil, a secondary structure composition that corresponds well both to the CD analysis of the bacterial aminopeptidase which is 35% α -helix, 17% β -sheet, and 48% coil [23] and the secondary structure composition of the model (33% α -helix, 24% β -sheet, and 43% coil). Not only does the proportion of each secondary structure type correspond well between the QC model and the bacterial aminopeptidase family, but also the predicted linear arrangement and position of each secondary structural element from Pspired corresponds well with the aminopeptidases (Figure 2). In addition, the model predicts that the conserved cysteine residues are in position to form a disulfide bond,

a prediction experimentally confirmed in our previous study [22]. Further, both N-linked glycosylation sites are located on the exterior of the protein model.

In contrast with human QC, papaya QC is an all β -sheet protein. This has been shown by Oberg [20], based on CD spectra. The computational methods used here are in agreement with this result, in that β -structure is predicted by Psipred and the weak similarity to a β -propeller structure is also suggestive of a β -structure. In addition, the lack of sequence similarity between the two enzymes [15], and the dramatic differences in susceptibility to inhibition by both competitive inhibitors and diethylpyrocarbonate [10,24] suggest that the two enzymes are unlikely to be divergently evolved.

Convergently evolved functions and/or mechanisms are not without precedent. The most widely known example is the catalytic triad, which was first found in protease chymotrypsin [25], and subsequently in the structurally unrelated but functionally similar subtilisin [26]. It has since been observed in lipases [27] and recently in acetylhydrolase [28]. All of these enzymes share related functions and similar mechanisms, but have quite different folds.

Holbrook and Ingram [29] have suggested a set of tests for distinguishing divergent and convergent evolution in enzymes. In descending order of importance they are: sequence similarity at the DNA level, sequence similarity at the amino-acid level, similarity of three-dimensional structure, similarity of enzyme-substrate interactions, similar catalytic mechanism and segments of the chain essential for catalysis are in the same sequence. The catalytic triad containing enzymes fail every test except for similar mechanism and so are generally accepted as examples of convergent evolution. Similar function with different fold and lack of sequence similarity is currently used as evidence for convergence (for example [30]). Russell [31] has looked for similar spatial arrangements of catalytic residues in proteins that share no fold similarity, and therefore are unlikely to be divergently evolved. In a search of the protein structure database he identified 22 pairs of proteins that have similar function but different folds, and are therefore likely to be convergently evolved. Similarly, we find that human and papaya QCs are unrelated at the DNA and amino-acid levels, are likely to have different structures, and yet have similar functions. In the absence of more structural studies we are unable to tell whether the least important of Holbrook and Ingram's criteria are met, but given the known structural differences this seems highly unlikely.

Six amino acid residues are predicted to be at the active site of QC based on their location in the zinc aminopepti-

dase fold (Figure 3, Table 1). These are picked up quickly in a PSI-BLAST [32] search with the human QC protein sequence. Two of these, QC histidines 140 and 330, were previously shown to be essential catalytic residues [10]. The other four residues, glutamates 201 and 202 and aspartates 159 and 248, were examined in this study. Mutagenesis of the two glutamates showed both to be essential for QC enzymatic activity. The aminopeptidase counterpart of QC Glu 201 acts as a general base to promote water attack on the peptide bond in aminopeptidase catalysis. A glutamate residue plays the same role in other well-studied zinc peptide hydrolases such as carboxypeptidase A and thermolysin. It is tempting to speculate that QC Glu 201 may perform a similar function in the transamidation reaction of glutaminy cyclization by promoting attack of the alpha amine on the gamma carboxamide.

Does human QC contain active site zinc? Our ICP-MS analysis indicated a stoichiometry of 0.3 mol Zn/mol QC, effectively ruling out the presence of two zinc atoms and suggesting the residual zinc presence is adventitious, although admittedly it is possible some zinc was lost during enzyme purification. QC is refractory to metal chelators such as EDTA, nitrolotri-acetic acid [5,10], peptide thiols and boronic acids [24], Glu-Trp, and gamma-glutamate hydroxamate (unpublished results). In addition, the enzyme refolds into an active form after denaturation with urea [6,9]. Finally, an absence of metal is consistent with the observation that mutation of the two aspartates, which are zinc ligands in the aminopeptidase, had no effect on QC enzymatic activity. Taken together, the above evidence argues strongly against a requirement for zinc at the QC active site.

Schilling et al [24] have recently proposed that QC is a metalloenzyme based on time dependent inhibition by o-phenanthroline and dipicolinic acid, as well as the activation of the inactive QC "apoenzyme" by added zinc. In contrast, we have observed competitive inhibition (K_i 20 μ M) with o-phenanthroline. As Schilling *et al.* point out, both o-phenanthroline and dipicolinic acid are planar heterocyclic compounds. Both can therefore potentially act as reversible inhibitors just as the imidazole derivatives they report [5]. It is also curious that cobalt did not reactivate QC in the Schilling study, since it commonly reactivates zinc peptidases. It remains to be seen, therefore, whether the reported reactivation by zinc is due to replacement of an active site metal, binding of zinc to an alternative site such as the polyhistidine expression tag, or an experimental artifact.

The bacterial zinc aminopeptidase fold is found in proteins such as carboxypeptidase G2 [19] as well as the human transferrin receptor [3], which exhibits no known

Table 1: Active site residue comparison. Comparison of positionally identical active site residues in *Vibrio proteolyticus* aminopeptidase and human pituitary glutaminyl cyclase. A designation of "none" means the residue plays no discernable role in the enzymatic reaction.

Aminopeptidase Residue	Function in Aminopeptidase	QC residue	Function in QC
Asp 117	Zn bridge	Asp 159	None
His 256	Coordinates Zn #1	His 330	Catalytic
Glu 152	Coordinates Zn #1	Glu 202	Catalytic
His 97	Coordinates Zn #2	His 140	Catalytic
Asp 179	Coordinates Zn #2	Asp 248	None
Glu 151	Catalytic	Glu 201	Catalytic
His 230	None	His 307	Substrate binding
Met 242	None	His 319	Substrate binding

enzymatic activity. A VAST search of the protein databank using the *Vibrio proteolyticus* aminopeptidase (1AMP) structure reveals a number of similar tertiary structures including, intriguingly, a pyroglutamate aminopeptidase (1A2Z) from *Thermococcus litoralis*. The VAST superimposition of 1A2Z with 1AMP shows good structural alignment with the central beta sheet and connecting helices, with an alpha carbon RMSD of 3.3 angstroms. Pyroglutamate aminopeptidase is a cysteine peptidase, with the catalytic triad only slightly offset from the zinc aminopeptidase active site residues described in this paper. It is clear, then, that the bacterial zinc aminopeptidase fold is available for a variety of functions other than zinc-catalyzed peptide bond hydrolysis. In the case of QC, it appears that four of the six critical active site residues of the aminopeptidase are also essential for QC catalysis, possibly representing the distant recruitment of catalytic residues during evolution among hydrolases/amidotransferases that operate on peptide termini.

Conclusions

A theoretical model of human pituitary glutaminyl cyclase has been generated based on the tertiary structure of the bacterial zinc aminopeptidase family. Six key active site residues in the aminopeptidase structure are present in the QC model, with four of these six residues essential for QC enzymatic activity. Despite the structural similarity with the zinc aminopeptidase family, our site directed mutagenesis, inhibitor studies, and metal analysis argue strongly against QC being a metalloenzyme.

The structural model proposed here will provide the basis for interpreting further mutagenic and spectroscopic studies of this mammalian peptide processing enzyme, as well as facilitate structural comparisons with the plant glutaminyl cyclase family.

Methods

Materials

Peptide substrates were obtained from Bachem. All restriction enzymes and their corresponding buffers were obtained from Promega. Oligonucleotide primers were made by Integrated DNA technologies, Inc. Automated sequencing was performed at UF DNA Core Lab. Mutagenesis reagents were obtained from Amersham Pharmacia Biotech. Human pituitary glutaminyl cyclase cDNA was expressed in *Drosophila* cell culture and purified from serum-free media by affinity chromatography as described previously [22].

Comparative modeling

Secondary structure predictions were calculated using Pspired [33]. Homologous proteins of known structure were identified, and initial sequence alignments were produced using the Fugue program [17]. Clear homologues were found: aminopeptidase from *Vibrio proteolyticus* (pdb code 1AMP) and *Streptomyces griseus* (pdb code 1XJO), and carboxypeptidase G2 from *Pseudomonas* sp. (pdb code 1CG2). These all belong to the "bacterial aminopeptidase" family in the structural alignment database Homstrad [34]. The initial alignment was adjusted manually (Figure 2) and verified using Harmony3 (J. Shi, unpublished), which matches observed substitution frequency in the alignment with predicted substitution frequency from standard and environment specific substitution tables. Models were built with Modeller 6.1 [35,36]. Since Cys 139 and Cys 164 were within bonding distance in the initial models, in subsequent models they were built in the disulfide bonded rather than free thiol form. Five loops (residues 177–186, 203–215, 227–237, 273–269, 298–304) contained insertions in the QC model relative to the template structures. These loops were rebuilt with Rapper [37,38]. As these loops had no equivalent in the parent structures, they should be considered the least reliable parts of the model. Similarly, no homologous residues were available for the N-terminal 35 residues and these were therefore omitted. The final model had only a

single Ramachandran outlier (Phe 146), according to the Rampage program [39]. It has been submitted to the Protein Data Bank and assigned pdb code 1MOI.

Site-directed mutagenesis

A mutation was introduced into the DES hQC expression vector using Unique Site Elimination Mutagenesis kit (Amersham Pharmacia Biotech). This method uses two primers to incorporate mutations into the same strand of DNA by in vitro DNA synthesis with T4 DNA polymerase [40]. Primers were designed to replace the glutamic acids at positions 201 or 202, and aspartic acids at positions 159 and 248 with alanine while introducing a silent mutation in the expression vector that deletes a restriction site. Mutant constructs were confirmed by automated sequencing at UF DNA Core Lab (Gainesville, FL).

Activity assay

A spectrophotometric assay was used to detect QC activity as reported by Bateman [41] with the modification that 5 mM Gln-NH₂ was routinely used as substrate and 10 μM bestatin was included. For kinetic analysis, substrate concentrations from 0.3 mM to 4 mM were used.

Protein samples (1–5 μg) were run in 10 % NuPAGE as described in the manufacturers instructions except the reducing agent, 50 μM DTT, was omitted in non-reduced samples.

Metal analysis

Purified protein samples were analyzed for metal content with ICP-MS by the University of Georgia Chemical Analysis Laboratory. Two independent purifications were used, with the buffers treated with Chelex 100 to remove residual zinc. Blanks included buffer alone as well as elution buffer from a complete purification protocol using media from nontransfected S2 cells which had been treated in a manner identical to the QC-expressing S2 cells. Protein was quantified using the predicted molar extinction coefficient (55 310 M⁻¹cm⁻¹ at 280 nm) and by SDS-PAGE followed by staining with Coomassie blue, then densitometry with BSA standards. Both methods agreed closely.

Secondary structure analysis

Purified protein samples were dialyzed against 5 mM potassium phosphate buffer, pH 6.0 and then concentrated to 6 μM using Microcon Centricon-10 (Amicon). Data was collected using an OLIS circular dichroism (CD) spectrophotometer. Fifty data points were collected for each scan from 196–260 nm with a total of eight scans per sample. Baseline corrected data were smoothed with a Gavitsky-Golay filter. The Y-scale was measured in ellipticity, θ, and set from -40 to 60. Molar ellipticity, [θ], was calculated for each data point using the following for-

mula: $[\theta] = (100 * \theta) / (C * n * l)$ where C = concentration in mM, n = number of residues, and l = path length of cell in cm. The molar ellipticity values were imported into the K2d program [42] which uses a database comparison to predict the % α-helix, % β-sheet, and % random coil.

List of abbreviations

QC, Glutaminyl Cyclase

ICP-MS, Inductively Coupled Plasma Mass Spectrometry

DES, Drosophila Expression System

PDB, Protein Data Bank

DTT, Dithiothreitol

SDS-PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis

BSA, bovine serum albumin

Authors' contributions

REB carried out the expression, purification, mutagenesis, and spectral studies. SCL performed the modeling studies. SAM performed the kinetic experiments. RCB designed and oversaw the study. All authors read and approved the final manuscript.

Acknowledgements

Funding from the NSF EPSCOR Program, the state of Mississippi, and the University of Southern Mississippi is gratefully acknowledged. SCL is a Wellcome Trust fellow of mathematical biology. We thank Iris Lindberg and Jae Hwang at the LSU Medical Center for advice on the S2 cell cultures. We also thank Pernilla Wittung-Stafshede and Irina Pozdnyakova at Tulane for assistance with the circular dichroism experiments.

References

- Steiner DF: **The proprotein convertases.** *Curr Opin Chem Biol* 1998, **2**:31-39.
- Reznik SE, Fricker LD: **Carboxypeptidases from A to z: implications in embryonic development and Wnt binding.** *Cell Mol Life Sci* 2001, **58**:1790-1804.
- Prigge ST, Mains RE, Eipper BA, Amzel LM: **New insights into copper monooxygenases and peptide amidation: structure, mechanism and function.** *Cell Mol Life Sci* 2000, **57**:1236-1259.
- Messer M, Ottesen M: **Isolation and properties of glutamine cyclotransferase of dried papaya latex.** *C R Trav Lab Carlsberg* 1965, **35**:1-24.
- Busby WH Jr, Quackenbush GE, Humm J, Youngblood WW, Kizer JS: **An enzyme that converts glutaminyl-peptides into pyroglutaminyl-peptides.** *J Biol Chem* 1987, **262**:8532-8536.
- Fischer WH, Spiess J: **Identification of a mammalian glutaminyl cyclase converting glutaminyl into pyroglutaminyl peptides.** *Proc Natl Acad Sci U S A* 1987, **84**:3628-3632.
- Bockers TM, Kreutz MR, Pohl T: **Glutaminyl-cyclase expression in the bovine/porcine hypothalamus and pituitary.** *J Neuroendocrinol* 1995, **7**:445-453.
- Pohl T, Zimmer M, Mugele K, Spiess J: **Primary structure and functional expression of a glutaminyl cyclase.** *Proc Natl Acad Sci U S A* 1991, **88**:10059-10063.

9. Song I, Chuang CZ, Bateman RC Jr: **Molecular cloning, sequence analysis and expression of human pituitary glutaminyl cyclase.** *J Mol Endocrinol* 1994, **13**:77-86.
10. Bateman RC Jr, Temple JS, Misquitta SA, Booth RE: **Evidence for essential histidines in human pituitary glutaminyl cyclase.** *Biochemistry* 2001, **40**:11246-11250.
11. Schilling S, Hoffmann T, Rosche F, Manhart S, Wasternack C, Demuth HU: **Heterologous expression and characterization of human glutaminyl cyclase: evidence for a disulfide bond with importance for catalytic activity.** *Biochemistry* 2002, **41**:10849-10857.
12. Temple JS, Song I, Burns KH, Bateman RC Jr: **Absence of an Essential Thiol in Human Glutaminyl Cyclase: Implications for Mechanism.** *Korean J Biol Sci* 1998, **2**:243-248.
13. Gololobov MY, Wang W, Bateman RC Jr: **Substrate and inhibitor specificity of glutamine cyclotransferase (QC).** *Biol Chem Hoppe Seyler* 1996, **377**:395-398.
14. Gololobov MY, Song I, Wang W, Bateman RC Jr: **Steady-state kinetics of glutamine cyclotransferase.** *Arch Biochem Biophys* 1994, **309**:300-307.
15. Dahl SW, Slaughter C, Lauritzen C, Bateman RC Jr, Connerton I, Pedersen J: **Carica papaya glutamine cyclotransferase belongs to a novel plant enzyme subfamily: cloning and characterization of the recombinant enzyme.** *Protein Expr Purif* 2000, **20**:27-36.
16. Chevrier B, Schalk C, D'Orchymont H, Rondeau JM, Moras D, Tarnus C: **Crystal structure of *Aeromonas proteolytica* aminopeptidase: a prototypical member of the co-catalytic zinc enzyme family.** *Structure* 1994, **2**:283-291.
17. Shi J, Blundell TL, Mizuguchi K: **FUGUE: sequence-structure homology recognition using environment-specific substitution tables and structure-dependent gap penalties.** *J Mol Biol* 2001, **310**:243-257.
18. Greenblatt HM, Almog O, Maras B, Spungin-Bialik A, Barra D, Blumberg S, Shoham G: **Streptomyces griseus aminopeptidase: X-ray crystallographic structure at 1.75 Å resolution.** *J Mol Biol* 1997, **265**:620-636.
19. Rowsell S, Paupit RA, Tucker AD, Melton RG, Blow DM, Brick P: **Crystal structure of carboxypeptidase G2, a bacterial enzyme with applications in cancer therapy.** *Structure* 1997, **5**:337-347.
20. Oberg KA, Ruysschaert JM, Azarkan M, Smolders N, Zerhouni S, Wintjens R, Amrani A, Looze Y: **Papaya glutamine cyclase, a plant enzyme highly resistant to proteolysis, adopts an all-beta conformation.** *Eur J Biochem* 1998, **258**:214-222.
21. Williams MG, Shirai H, Shi J, Nagendra HG, Mueller J, Mizuguchi K, Miguel RN, Lovell SC, Innis CA, Deane CM, Chen L, Campillo N, Burke DF, Blundell TL, de Bakker PI: **Sequence-structure homology recognition by iterative alignment refinement and comparative modeling.** *Proteins* 2001, **Suppl 5**:92-97.
22. Booth RE, Misquitta SA, Bateman RC Jr: **Human pituitary glutaminyl cyclase: expression in insect cells and dye affinity purification.** *Protein Expr Purif* 2003, **32**:141-146.
23. Ustynyuk L, Bennett B, Edwards T, Holz RC: **Inhibition of the aminopeptidase from *Aeromonas proteolytica* by aliphatic alcohols. Characterization of the hydrophobic substrate recognition site.** *Biochemistry* 1999, **38**:11433-11439.
24. Schilling S, Niestroj AJ, Rahfeld JU, Hoffmann T, Wermann M, Zunkel K, Wasternack C, Demuth HU: **Identification of human glutaminyl cyclase as a metalloenzyme: Potent inhibition by imidazole derivatives and heterocyclic chelators.** *J Biol Chem* 2003.
25. Blow DM: **The study of α -chymotrypsin by x-ray diffraction. The third Ciba medal lecture.** *Biochemical Journal* 1969, **112**:261-268.
26. Alden RA, Birktoft JJ, Kraut J, Robertus JD, Wright CS: **Atomic coordinates for subtilisin BPN' (or Novo).** *Biochem Biophys Res Commun* 1971, **45**:337-344.
27. Brady L, Brzozowski AM, Derewenda ZS, Dodson E, Dodson G, Tolley S, Turkmenburg JP, Christiansen L, Høge-Jensen B, Nørskov L: **A serine protease triad forms the catalytic centre of a triacylglycerol lipase.** *Nature* 1990, **343**:767-770.
28. Ho YS, Swenson L, Derewenda U, Serre L, Wei Y, Dauter Z, Hattori M, Adachi T, Aoki J, Arai H, Inoue K, Derewenda ZS: **Brain acetylhydrolase that inactivates platelet-activating factor is a G-protein-like trimer.** *Nature* 1997, **385**:89-93.
29. Holbrook JJ, Ingram VA: **Ionic properties of an essential histidine residue in pig heart lactate dehydrogenase.** *Biochem J* 1973, **131**:729-738.
30. Beuth B, Niefind K, Schomburg D: **Crystal structure of creatinase from *Pseudomonas putida*: a novel fold and a case of convergent evolution.** *J Mol Biol* 2003, **332**:287-301.
31. Russell RB: **Detection of protein three-dimensional side-chain patterns: new examples of convergent evolution.** *J Mol Biol* 1998, **279**:1211-1227.
32. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.** *Nucleic Acids Res* 1997, **25**:3389-3402.
33. McGuffin LJ, Bryson K, Jones DT: **The PSIPRED protein structure prediction server.** *Bioinformatics* 2000, **16**:404-405.
34. Mizuguchi K, Deane CM, Blundell TL, Overington JP: **HOMSTRAD: a database of protein structure alignments for homologous families.** *Protein Sci* 1998, **7**:2469-2471.
35. Sali A, Blundell TL: **Comparative protein modelling by satisfaction of spatial restraints.** *J Mol Biol* 1993, **234**:779-815.
36. Fiser A, Do RK, Sali A: **Modeling of loops in protein structures.** *Protein Sci* 2000, **9**:1753-1773.
37. de Bakker PI, DePristo MA, Burke DF, Blundell TL: **Ab initio construction of polypeptide fragments: Accuracy of loop decoy discrimination by an all-atom statistical potential and the AMBER force field with the Generalized Born solvation model.** *Proteins* 2003, **51**:21-40.
38. DePristo MA, de Bakker PI, Lovell SC, Blundell TL: **Ab initio construction of polypeptide fragments: efficient generation of accurate, representative ensembles.** *Proteins* 2003, **51**:41-55.
39. Lovell SC, Davis IW, Arendall WB III, de Bakker PI, Word JM, Prisant MG, Richardson JS, Richardson DC: **Structure validation by Calpha geometry: phi, psi and Cbeta deviation.** *Proteins* 2003, **50**:437-450.
40. Ray FA, Nickoloff JA: **Site-specific mutagenesis of almost any plasmid using a PCR-based version of unique site elimination.** *Biotechniques* 1992, **13**:342-348.
41. Bateman RC Jr: **A spectrophotometric assay for glutaminyl-peptide cyclizing enzymes.** *J Neurosci Methods* 1989, **30**:23-28.
42. Andrade MA, Chacon P, Merelo JJ, Moran F: **Evaluation of secondary structure of proteins from UV circular dichroism spectra using an unsupervised learning neural network.** *Protein Eng* 1993, **6**:383-390.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

